



Saskatchewan

Invasive Plant Species Identification Guide



LEGEND

Plant Shape



Rosette



Branched



Rosette in seedling year



Slightly branched



Multi-stemmed



Not branched



Shrub or tree



Low & spreading



Highly branched



Narrow with drooping seed head

Invasive Plant Species Identification Guide

Introduction

This guide is to be used as a tool to help identify invasive plant species in Saskatchewan. An invasive plant species is a non-native plant that, because of its growth and reproductive characteristics, can adapt to one or more habitats very quickly and proliferate, expanding in range. An invasive plant species has a negative economic, ecological, and/or aesthetic effect.

The Invasive Plant Species Identification Guide for Saskatchewan has been designed for agricultural producers, land managers, and visitors to Saskatchewan. Producers and land managers play an important role in invasive species management as they are on the ground everyday, acutely aware of changes on the landscape. Visitors to the area often have a keen eye for the vegetation they

are seeing. Visitors need to be aware of the threat invasive species pose so that these plants are not brought into, spread around or transported out of the area.

Invasive species are a growing threat to the integrity of native prairie. Native prairie is home to many important and/or rare plants and animals. The plants which make up native prairie work together to store carbon, maintain water and nutrient cycling, and build soil. Invasive species can upset this balance.

The invasion of non-native species in haylands, tame forage, gravel pits, riparian areas, roadsides and cropland affect the function of these habitats and can cause a downgrade in their usefulness.

All of the species included in this guide are a threat to Saskatchewan.

Prevention of invasive species is the most cost effective and environmentally responsible management tool. Do not transport hay, soil, or plants from an area with known infestations to an uninfested area. Most invasive species require some bare ground to establish. Keeping permanent cover in a healthy range condition minimizes the amount of habitat an invasive species can colonize. Control new infestations in cultivated fields when they are first noticed. Wash or check clothing, pets, horses, livestock, equipment and vehicles for plant parts or seeds before coming into or leaving a new area.

Early identification increases the chances of eradication. Take the time to identify that plant that looks out of place, you've never seen before, or has suspiciously shown up. It could save you or your neighbours time and money in the long run!

Some of the invasive plants in this guide have not yet been identified in Saskatchewan, or have only been noted in very isolated populations. They have been included in this guide so that if they are present, a rapid identification and control can take place prior to an exponential invasion. If you identify one of these invasive plants contact the Provincial Weed Control Specialist at 306-787-4673.

These not-yet locally identified or uncommon species include: Yellow star thistle, Field scabious, Purple loosestrife, Salt cedar, Spotted knapweed, Russian knapweed, Diffuse knapweed, African rue, Hoary cress/White top, and Orange hawkweed.

If you identify an invasive plant species consult with your local Rural Municipality, your local Weed Management Area, a certified Agronomist or Agrologist, or the Saskatchewan Ministry of Agriculture Agricultural Knowledge Centre at 1-866-457-2377.

Table of Contents

Plant Shapes	4
Diagrams of Plant Parts	5
Plant Species Listings	7
Grasses	
Downy brome - <i>Bromus tectorum</i>	8
Japanese brome - <i>Bromus japonicus</i>	9
Yellow Flowered Plants	
Absinthe wormwood - <i>Artemisia absinthium</i>	10
Common tansy - <i>Tanacetum vulgare</i>	11
Dalmatian toadflax - <i>Linaria dalmatica</i>	12
Yellow toadflax - <i>Linaria vulgaris</i>	13
Leafy spurge - <i>Euphorbia esula</i>	14
Yellow star thistle - <i>Centaurea solstitialis</i>	15
Purplish Flowered Plants	
Bull thistle - <i>Cirsium vulgare</i>	16
Canada thistle - <i>Cirsium arvense</i>	17
Nodding thistle - <i>Carduus nutans</i>	18
Field scabious - <i>Knautia arvensis</i>	19
Purple loosestrife - <i>Lythrum salicaria</i>	20
Dames Rocket - <i>Hesperis matronalis</i>	21
Pink Flowered Plants	
Common burdock - <i>Arctium minus</i> subsp. <i>minus</i>	22
Salt cedar - <i>Tamarix spp.</i>	23
Spotted knapweed - <i>Centaurea maculosa</i>	24
Russian knapweed - <i>Acroptilon repens</i>	25
White Flowered Plants	
Diffuse knapweed - <i>Centaurea diffusa</i>	26
African rue - <i>Peganum harmala</i>	27
Baby's breath - <i>Gypsophila paniculata</i>	28
Field bindweed - <i>Convolvulus arvensis</i>	29
Hoary cress or white top - <i>Cardaria draba</i>	30
Round-leaved mallow - <i>Malva pusilla</i> or <i>Malva rotundifolia</i>	31
Scentless chamomile - <i>Matricaria perforata</i>	32
Hoary alyssum - <i>Berteroa incana</i>	33
Ox-eye daisy - <i>Chrysanthemum leucanthemum</i>	34
Orange/Red Flowered Plants	
Orange hawkweed - <i>Hieracium aurantiacum</i>	35
Glossary	36

Plant Shapes



Rosette



Rosette in
seedling year



Multi-stemmed



Shrub or tree



Highly
branched



Branched



Slightly
branched



Not branched




Low &
spreading




Narrow with
drooping seed
head

The shape of a plant can be used as a distinguishing characteristic—setting it apart from other plants. Some plants can have more than one shape depending on their life cycle, habitat, or the climate.”


Leaf Parts




Grass Plant Parts



Shapes of Simple Leaves



Types of Divided Leaves



Plant species may be listed as noxious weeds in one or more jurisdictions when they are invasive and pose a threat of invasion, rapid spread and difficulty in control. It is important to be aware of invasive species in adjoining jurisdictions as the threat from neighbouring areas may be significant.

To determine the species listing of any invasive plant species in your area, please contact your local RM or the Saskatchewan Ministry of Agriculture



Downy Brome – *Bromus tectorum*

Also Known As: cheatgrass, thatch bromegrass, early chess, downy chess

General Description: hairy, purplish tinged winter or summer annual grass with fibrous roots; often growing in dense patches; seed spikelets droop to one side

Leaves: 0.75 to 6 inches (1.9 to 15.2 cm) long, green; soft hairs on the top and bottom of leaf; lacks auricles, but has a jagged 0.04 to 0.12 inch (0.1 to 0.3 mm) long hairless ligule

Stems: slender, upright, hairy; leaf sheath has long hairs; often purplish tinged

Height: 3 to 24 inches (7.6 to 61 cm) tall

Flower: drooping, spreading spikelets (of 4 or more seeds) on slender twisting branches; droops to one side; rough 0.38 to 0.75 inch (1.0 to 1.9 cm) long straight to slightly bent awns; purplish tinged

Habitat: tame forage, native prairie, yard sites, cropland, waste areas & roadsides

Time of Flowering: heads out in May through June with seed set shortly thereafter

Key Distinguishing Characteristic(s): similar to Japanese brome (*Bromus japonicus*) - distinguishable by smaller seeds, lack of silky hairs on the ligule, straight to slightly bent awns and a more spreading appearance of the drooping seed heads

Impact: displaces native vegetation, reduces crop yields and increases risks of wildfire; awns can cause injury to grazing livestock

Plant Shape





Japanese Brome - *Bromus japonicus*

Also Known As: Japanese chess, field brome

General Description: hairy, purplish tinged, winter or summer annual grass with fibrous roots; often growing in dense patches; drooping, compact, flattened seed spikelets

Leaves: 1.5 to 6 inches (3.8 to 15.2 cm) long, green; soft hairs on the top and bottom of leaf; lacks auricles, but has a jagged 0.02 to 0.08 inches (0.05 to 0.2 cm) long hairy ligule

Stems: slender, upright, hairy; leaf sheath has long hairs

Height: 3 to 27 inches (7.6 to 69 cm) tall

Flower: slightly drooping spikelets (of 4 or more seeds) on slender twisting branches; rough 0.38 to 0.75 inch (1.0 to 1.9 cm) long sharply bent and/or twisted awns; purplish tinged

Habitat: tame forage, native prairie, yard sites, cropland, waste areas & roadsides

Time of Flowering: heads out in May through June with seed set shortly thereafter

Key Distinguishing Characteristic(s): similar to downy brome (*Bromus tectorum*)- distinguishable by silky hairs on the ligule, compact appearance of slightly drooped seed heads, and sharply bent and/or twisted awns

Impact: displaces native vegetation, reduces crop yields and increases risks of wildfire; awns can cause injury to grazing livestock

Plant Shape





Absinthe Wormwood – *Artemisia absinthium*

Also Known As: absinthe, wormwood, wormwood sage

General Description: greenish-grey, multi-stemmed perennial with tap root growth; top growth dies back yearly

Leaves: 1 to 3 inches (2.5 to 7.6 cm) long, greenish-grey, silky haired; divided into deep segments or lobes; on long stalks near base- shorter higher up the stem; alternate on the stem

Stems: woolly, grooved, greenish-grey; many stems come from basal rosette

Height: up to 60 inches (152 cm) tall

Flower: small yellowish tubular flowers in clusters; flower clusters 0.08 to 0.12 inches (0.2 to 0.3 cm) in diameter; inconspicuous and plentiful

Habitat: tame forage, waste areas, yard sites, cropland, roadsides & gravel pits

Time of Flowering: late July through September

Key Distinguishing Characteristic(s): strong sage-like odour; divided and silky haired leaves on stems from a rosette

Impact: competes with other vegetation; taints hay (not palatable to livestock)

Plant Shape





Common Tansy – *Tanacetum vulgare*

Also Known As: garden tansy, yellow buttons

General Description: aromatic, perennial with tap and creeping root system; rhizome root growth; distinct flat topped, yellow button-like flowers

Leaves: 4 to 8 inch (10.2 to 20.3 cm) long, green, almost fern-like; on short stalks, with jagged, deeply lobed segments; strong smelling; alternate on the stem

Stems: slightly branched, often purplish-red, dotted with small glands; many branches grow from the same rootstock

Height: 18 to 36 inches (46 to 91 cm) tall

Flower: yellow, daisy-like without petals; grouped on the tops of the plants

Habitat: riparian areas, tame forage, native prairie, roadsides, waste areas & gravel pits

Time of Flowering: July through September

Key Distinguishing Characteristic(s): Button-like flowers (daisy without petals) and sharply toothed leaves

Impact: chokes out other vegetation; although generally avoided by livestock, if consumed, it can have toxic effects ranging from none to abortions and in very rare cases death

Plant Shape





Dalmatian Toadflax – *Linaria dalmatica*

Also Known As: butter and eggs, wild snapdragon, *Linaria genistifolia* spp. *dalmatica*

General Description: perennial with tap root and creeping root system; snapdragon-like flower heads; produces clear to slightly milky juice when leaves or stems are broken

Leaves: 0.75 to 2.4 inches (1.9 to 6.0 cm) long, waxy green, hairless; heart-shaped and clasping the stem; alternate on the stem

Stems: slightly branched with woody base and thick waxy coating

Height: 15 to 48 inches (38 to 122 cm) tall

Flower: light yellow; snapdragon-like with prominent spur at the flower base; located along the upper portion of the stems in a spike like cluster

Habitat: tame forage, native prairie, cropland, yard sites, roadsides & gravel pits

Time of Flowering: late June through August, possibly into September

Key Distinguishing Characteristic(s): broad, heart-shaped leaves which clasp the stem; snapdragon-like yellow flowers; grows in diffuse patches

Impact: chokes out other vegetation; can hybridize with yellow toadflax

Plant Shape





Yellow Toadflax – *Linaria vulgaris*

Also Known As: butter and eggs, Jacob's ladder, wild snapdragon

General Description: perennial with tap root and creeping root systems; snapdragon-like flower heads; produces clear to slightly milky juice when leaves or stems are broken

Leaves: 0.75 to 4 inches (1.9 to 10.2 cm) long; pale to silvery-green (waxy), hairless; long and lance-shaped on short leaf stems; alternate, but can appear almost opposite when crowded on the stem

Stems: hairless and woody red at base; slightly hairy and green near top

Height: 4 to 36 inches (10.2 to 91 cm) tall

Flower: light yellow with orange centers; snapdragon-like with prominent spur at the flower base; in spike-like clusters along the upper portion of the stems

Habitat: tame forage, native prairie, cropland, yard sites, roadsides & gravel pits

Time of Flowering: late June through August, possibly into September

Key Distinguishing Characteristic(s): long, narrow, alternate leaves on short leaf stems; snapdragon-like flowers; grows in dense patches

Impact: chokes out other vegetation and is slightly toxic to livestock; can hybridize with dalmatian toadflax

Plant Shape





Leafy Spurge – *Euphorbia esula*

General Description: perennial with tap root and creeping root system; single stem plant grown in a cluster; distinct yellowish-green bracts emerge in June on plant tops

Leaves: 0.25 to 1.5 inches (0.6 to 3.8 cm) long, green, hairless; long and lance-shaped; alternate on the stem

Stems: hairless, pale green to blue-green

Height: 6 to 36 inches (15 to 91 cm) tall

Flower: inconspicuous, yellowish-green; small, less than 0.12 inches (0.3 cm) wide, lacks petals; flowers sit within clustered, showy yellowish-green bracts on plant tops

Habitat: riparian areas, tame forage, native prairie, coulees, cropland, roadsides, yard sites, waste areas & gravel pits

Time of Flowering: bracts appear in early June; true flowers appear end of June and continue to flower into the fall

Key Distinguishing Characteristic(s): milky latex sap is produced when stems or leaves are broken

Impact: chokes out other vegetation; can be toxic to cattle and horses

Plant Shape





Yellow Star Thistle – *Centaurea solstitialis*

General Description: winter annual with deep tap root; yellow flower head surrounded by a star of long sharp thorns

Leaves: basal leaves generally 2 to 3 inches (5.0 to 7.6 cm) long and deeply lobed; long and narrow shaped upper leaves attach to wings running down the sides of stem

Stems: stiff, upright stems are covered in fine woolly silvery hair; grey-green to bluish-green

Height: 2 to 72 inches (5 to 183 cm) tall, with an average of 12 to 36 inches (30 to 61 cm) tall

Flower: yellow, tubular; flower heads sit on stem tips within bracts with sharp 0.25 to 2.0 inch (0.64 to 5.0 cm) long golden-coloured thorns arranged around the flower head in a star shape

Habitat: tame forage, native prairie, cropland, waste areas, roadsides & gravel pits

Time of Flowering: late June through August

Key Distinguishing Characteristic(s): star shape arrangement of sharp golden-coloured thorns on flower bracts

Impact: chokes out desired vegetation; high water consumer; thorns can injure livestock and wildlife; fatally toxic to horses if consumed

Plant Shape





Bull Thistle – *Cirsium vulgare*

General Description: biennial thistle with tap root growth; extremely spiny

Leaves: spiny upper surface and edges; hairy lower surface; deeply lobed with long spines on the ends of the lobes; alternate on the stem

Stems: slightly woolly; spiny wings cover the stems

Height: 12 to 60 inches (30 to 152 cm) tall; in the first year only forms a large flat rosette

Flower: rose to purple-coloured, rarely white; tubular; flower heads up to 3 inches (7.6 cm) in diameter and sit within hairy, yellowish-tipped, prickly bracts on the ends of branches

Habitat: tame forage, roadsides, abandoned cropland, native prairie, waste areas & gravel pits

Time of Flowering: throughout the summer

Key Distinguishing Characteristic(s): only thistle with spines on the leaf surface, not just on the leaf edges; stems winged with long pointed spines

Impact: chokes out other vegetation, but it does not withstand cultivation

Plant Shape





Canada Thistle – *Cirsium arvense*

Also Known As: California thistle, field thistle, creeping thistle

General Description: perennial thistle with tap root and extensive creeping root system; grows in patches which can spread quickly; plants are either male or female

Leaves: numerous, light to dark green, hairless to cobwebby undersides; long with irregular shaped spine-toothed lobes; alternate on the stem

Stems: slightly prickly, wingless; light to dark green

Height: up to 48 inches (122 cm) tall

Flower: rose-purple to pink-white, tubular; flower heads numerous, up to 0.75 inches (1.9 cm) wide; male (globe-shaped) and female (flask-shaped) flower heads on separate plants; sits within almost spineless bracts on the ends of branches

Habitat: riparian areas, tame forage, native prairie, fence rows, waste areas, yard sites, cropland, roadsides & gravel pits

Time of Flowering: throughout summer

Key Distinguishing Characteristic(s): small flower heads and green, wingless but prickly stems; spines only on leaf edges; male and female flowers are found on different plants

Impact: chokes out other vegetation; new plants can grow from root segments

Plant Shape





Nodding Thistle – *Carduus nutans*

Also Known As: musk thistle, plumeless thistle

General Description: biennial with long tap root; nodding flower heads on top of slightly branched plant

Leaves: up to 16 inches (41 cm) long, dark green, with deep, light green midrib and spiny toothed edges; deeply lobed in an irregular fashion; alternate on the stem

Stems: small spiny wings along the stem; very few branches; stems naked directly below the flower heads

Height: 12 to 72 inches (30 to 183 cm) tall

Flower: tubular, purple; 1 to 2.5 inch (2.5 to 6.4 cm) wide flower head; flower heads sit within spine-tipped bracts on branch tops; outer bracts bent at right angles to the flower head; flower heads often nod over

Habitat: tame forage, native prairie, roadsides, waste areas & gravel pits

Time of Flowering: July through September in the 2nd year of growth

Key Distinguishing Characteristic(s): nodding flower heads; outer bracts bent at right angles to the flower heads; naked stems directly below the flower heads

Impact: chokes out other vegetation

Plant Shape





Field Scabious – *Knautia arvensis*

Also Known As: blue buttons

General Description: single-stemmed perennial with tap root growth; pale blue to purple, half-sphere shaped flower heads

Leaves: hairy, 4 to 10 inches (10.2 to 25 cm) long; rosette leaves are jaggedly toothed; stem leaves are deeply lobed and opposite on the stem

Stems: tall, slightly branched with stiff hairs covering the stem

Height: up to 60 inches (152 cm) tall

Flower: small, pale blue to purple coloured; clustered into 1 to 1.5 inch (2.5 to 3.8 cm) wide, half-sphere shaped flower heads on the ends of long leafless branches; very few flower heads per plant

Habitat: riparian areas, tame forage, waste areas, native prairie, abandoned cropland, yard sites, cropland, roadsides & gravel pits

Time of Flowering: June through early fall

Key Distinguishing Characteristic(s): tall, slightly branched with hairy leaves and stems; pale blue to purple coloured, half-sphere shaped flower heads

Impact: competes with other vegetation; not palatable to livestock and wildlife

Plant Shape





Purple loosestrife – *Lythrum salicaria*

Also Known As: lythrum, purple lythrum

General Description: wetland perennial; extensive tap root and creeping root system sprouts many square-shaped stems which support loose spikes of magenta-coloured flowers

Leaves: dark green, opposite or whorled in 3's, attached directly to the stem; narrow with a rounded base and smooth leaf edges

Stems: square-shaped (sometimes 5 or 6-sided); hairless to slightly hairy; one plant can sprout more than 30 stems

Height: 18 to 78 inches (46 to 198 cm) tall

Flower: 5 – 7 reddish-purple to magenta-coloured petals surrounding a small yellow center; loosely clustered into spikes along stem tops

Habitat: riparian areas & wetlands

Time of Flowering: July through September

Key Distinguishing Characteristic(s): square-shaped stems, magenta flowers and narrow leaves without stalks

Impact: chokes out desired wetland vegetation

Plant Shape





Dames Rocket – *Hesperis matronalis*

Also Known As: common dames violet, sweet rocket, mother-of-the-evening

General Description: dark green tall biennial or perennial with rosette in seedling year; single or multi-stemmed 2nd year growth with a tap root system; belongs to the mustard family; is a prolific seed producer

Leaves: 2 to 6 inches (5 to 15 cm) long, 0.4 to 1.6 inches (1 to 4 cm) wide, dark green, hairy on top and bottom; leaves get smaller further up the stem; jagged toothed edges; oblong to oblanceolate shaped; alternate on the stem

Stems: hairy; stiff & erect with some branching at the top; sometimes there is more than one stem per plant

Height: up to 39 inches (100 cm) tall; may be taller in riparian areas

Flower: 4 petals; fragrant flowers in loose clusters (like a loose ball) at the branch ends; usually purple in color but pink and white varieties exist

Habitat: riparian areas, gardens, shorelines, abandoned cropland, wooded areas, shrub bluffs, native prairie, coulees, yard sites, and waste areas

Time of Flowering: May through June or July

Key Distinguishing Characteristic(s): fragrant 4-petal flowers in loose clusters & alternate leaves

Impact: crowds out native vegetation because of large number of seeds released; decreases riparian area health

Plant Shape





Common Burdock – *Arctium minus* subsp. *minus*

Also Known As: wild rhubarb, beggar's button, elephant's ear, wild burdock, lesser burdock

General Description: tap rooted biennial; large alternate leaves

Leaves: somewhat heart shaped, similar to rhubarb leaves; deeply veined, alternate; up to 12 inches (30 cm) in length at the base, becoming smaller higher up the plant

Stems: tall, branched and hairy; stems and leaf stalks are hollow and grooved

Height: 24 to 72 inches (61 to 183 cm) tall at maturity

Flower: purple to white; disc-shaped flower heads at the ends of stems and at leaf axils along stems; 0.6 to 1.2 inches (1.5 to 3.0 cm) wide; heads surrounded by bracts with harsh, hooked spines

Habitat: riparian areas, coulees, yard sites, fencerows, waste areas & roadsides

Time of Flowering: mid-July through mid-September

Key Distinguishing Characteristic(s): tall stalky plant with very large alternate leaves (up to 12 inches or 30 cm long); large bur surrounds seeds

Impact: large burs impact livestock production and health

Plant Shape





Salt Cedar – *Tamarix* spp.

Also Known As: tamarisk

General Description: spindly perennial shrub or tree forming a dense thicket; small white to pink flowers cover the tree

Leaves: alternate, blue-green turning golden-orange in the fall before falling off; small (0.06 inches or 0.15 cm long), scale-like, lance-shaped; resemble juniper leaves

Stems: willowy, smooth, varying greatly in color when young; mature stems grey or brownish, grooved and wrinkled in appearance

Height: 60 to 240 inches (152 to 610 cm) tall at maturity

Flower: small, light pink to white; four to five 0.04 to 0.08 inch (0.1 to 0.2 cm) long petals; flowers can number in the thousands

Habitat: riparian areas & wetlands

Time of Flowering: spring through early fall

Key Distinguishing Characteristic(s): spindly tree with juniper-like leaves covered in small white to pink flowers

Impact: chokes out desired vegetation and concentrates salts near the soil surface to deter other species from growing; fire tolerant; high consumer of water (200 gal or 757 L/day)

Plant Shape





Spotted Knapweed – *Centaurea maculosa*

General Description: biennial or perennial with long tap root; pink to purple-coloured flowers

Leaves: somewhat hairy; deeply lobed dividing the leaf into narrow segments; alternate, usually clasping the stem; basal leaves “cobwebby” in appearance

Stems: branched, hairy; upper stems almost straight up and down

Height: 24 to 36 inches (61 to 91 cm) tall

Flower: usually pink to purple, tubular; flower heads 0.2 to 0.3 inches (0.6 to 0.8 cm) wide, numerous; flower heads sit on tops of leaf branches in oval bracts with a black fringe

Habitat: native prairie, tame forage, roadsides & waste areas

Time of Flowering: June through October

Key Distinguishing Characteristic(s): oval-shaped, black-fringed flower head bracts; “cobwebby” basal leaves; pink to purplish flowers

Impact: chokes out other vegetation; avoided by grazing animals

Plant Shape





Russian Knapweed – *Acroptilon repens*

Also Known As: *Centaurea repens*, hard heads

General Description: branched perennial with blackish, scaly, creeping roots; grows in dense patches

Leaves: narrow with irregular teeth at the base of the plant; upper leaves narrow, smooth edged with a spine at the tip; alternate, usually clasping the stem; woolly when young

Stems: highly branched; covered with soft grey hairs

Height: 24 to 36 inches (61 to 91 cm) tall

Flower: tubular flowers form a spherical flower head; flower heads 0.6 to 1.3 inches (1.5 to 3.3 cm) in diameter; sits on tops of leaf branches in entire bracts; silvery when unopened, pink to purple when open and straw-coloured when older

Habitat: cropland, native prairie, tame forage, roadsides & waste areas

Time of Flowering: July

Key Distinguishing Characteristic(s): upper leaves end in a soft spine but the rest of the plant generally lacks spines and prickles; flower head bracts are entire

Impact: chokes out other vegetation

Plant Shape





Diffuse Knapweed – *Centaurea diffusa*

Diffuse Knapweed

General Description: branched biennial or short-lived perennial with long tap root

Leaves: woolly, deeply lobed dividing the leaf into narrow segments; alternate leaves usually clasping the stem; basal leaves “cobwebby”

Stems: highly branched, hairy

Height: 24 to 36 inches (61 to 91 cm) tall

Flower: usually white to light purple, tubular; numerous; flower heads sit on stem tops in triangular, spine-tipped bracts

Habitat: native prairie, tame forage, roadsides & waste areas

Time of Flowering: July

Key Distinguishing Characteristic(s): triangular, spine-tipped flower head bracts; “cobwebby” basal leaves

Impact: chokes out other vegetation; avoided by grazing animals

Plant Shape





African Rue – *Peganum harmala*

Also Known As: harmel, harmal peganum, Syrian rue, isband

General Description: bright green shrubby perennial; branching roots to 20 feet deep; small white flowers and deeply cut leaves

Leaves: 1 to 2 inches (2.5 to 5.1 cm) long; deeply cut into long fine, narrow segments; bright green and alternate

Stems: fleshy, highly branched

Height: 12 to 24 inches (30 to 61 cm) tall at maturity

Flower: 5 white petals with yellow center; small, up to 1.0 inches (2.5 cm) in diameter; located along the stems in leaf axils

Habitat: native prairie, waste places, gravel pits & roadsides

Time of Flowering: June through August

Key Distinguishing Characteristic(s): bushy, succulent looking plant with small white flowers along the stems in the leaf axils

Impact: toxic to livestock; has allelopathic properties & replaces valuable vegetation

Plant Shape





Baby's Breath – *Gypsophila paniculata*

General Description: perennial with deep thick root; sparse foliage with open branched stems and numerous tiny flowers

Leaves: 1 to 4 inches (2.5 to 10.2 cm) long, blue-green to grey-green with glandular hairs; narrow, lance shaped with a single vein running up the middle of the leaf; opposite on the stem

Stems: slender, branched, blue-green to grey-green in colour; swollen where leaves attach to stems

Height: 12 to 36 inches (30 to 91 cm) tall

Flower: 5 white to rosy-coloured petals; tiny (less than 0.4 inches or 1.0 cm wide) and numerous on small spreading flower branches; gives the plant a “puff ball” appearance

Habitat: tame forage, native prairie, roadsides, waste areas & gravel pits

Time of Flowering: June through August; flowers for the first time in its 3rd year

Key Distinguishing Characteristic(s): highly branched; flowers and flower branches give the plant a large, round, sparse, airy “puff ball” appearance

Impact: crowds out other vegetation

Plant Shape





Field Bindweed – *Convolvulus arvensis*

Also Known As: creeping Jenny, European bindweed, morning glory, perennial morning glory

General Description: perennial with tap root and creeping root system; forms an extensive mat or climbs counterclockwise up nearby plants or objects

Leaves: dull green; arrow-head shaped, anchored by leaf stems off the main stem; numerous leaves of varied sizes on a single plant

Stems: smooth, 12 to 120 inches (30 to 305 cm) long; can wind counterclockwise around nearby plants or objects

Height: prostrate; needs to crawl up nearby objects and plants to achieve height

Flower: pinkish to white, up to 1 inch (2.5 cm) in diameter; morning glory or bell/trumpet shaped; on long flower stems with 2 small bracts part way down each flower stem

Habitat: cropland, riparian areas, tame forage, native prairie, waste areas, roadsides & gravel pits

Time of Flowering: June through September

Key Distinguishing Characteristic(s): 2 small bracts on the flower stems; climbs counterclockwise; large showy field bindweed flowers are easily distinguishable from wild buckwheat's small, inconspicuous flowers

Impact: chokes out other vegetation; makes harvesting crops and forages difficult

Plant Shape





Hoary Cress or White Top – *Cardaria draba*

Also Known As: pepper grass, white weed, heart-pod hoary cress

General Description: perennial, extensive creeping root system; small white flowers clustered on plant tops

Leaves: blue-green to grey-green; covered with fine hairs; egg to arrow-head-shaped; the large, narrow basal leaves have leaf stalks while the upper leaves have two lobes clasping the stem

Stems: greyish, hairy at base to hairless at the top of plant; slightly branched

Height: up to 24 inches (61 cm) tall

Flower: four small (0.12 inch or 0.3 cm long) white petals; flowers densely clustered on stem tops — appear flat topped

Habitat: roadsides, cropland, tame forage, native prairie, waste areas & gravel pits

Time of Flowering: May through July

Key Distinguishing Characteristic(s): clustered white flowers on tops of plants with large, finely haired leaves; seed pod is heart-shaped and contains two seeds

Impact: chokes out desired vegetation

Plant Shape





Round-Leaved Mallow – *Malva rotundifolia*

Also Known As: low mallow, *Malva pusilla*

General Description: annual to short-lived perennial; woody tap root; spreading stems with round to kidney-shaped leaves

Leaves: wrinkled, green, alternate; round to kidney-shaped with rounded or scalloped edges; on very long leaf stalks from the stem

Stems: prostrate, branched, up to 39 inches (100 cm) long

Height: spreads along ground with leaves reaching upwards

Flower: 1 to 3 small (0.2 inch or 0.5 cm), pale purple to white flowers grouped at the base of leaf stalks

Habitat: cropland, yard sites, tame forage, roadsides & waste areas

Time of Flowering: June through September

Key Distinguishing Characteristic(s): round to kidney-shaped leaves with scalloped edges on long leaf stalks

Impact: chokes out other vegetation; seeds stay viable for many years

Plant Shape





Scentless Chamomile – *Matricaria perforata*

Also Known As: scentless mayweed, scentless false mayweed

General Description: annual or biennial, rarely a perennial; dense fibrous roots; white daisy-like flowers

Leaves: smooth, very finely divided, almost carrot-like; alternate on the stem, usually without stalks; scentless

Stems: hairless, highly branched; upward curving branches

Height: 6 to 36 inches (15 to 91 cm) tall

Flower: daisy-like with yellow centre and white petals; 0.8 to 1.6 inches (2.0 to 4.0 cm) wide; occurs individually on stem tips

Habitat: roadsides, yard sites, riparian areas, cropland, tame forage, native prairie, waste areas & gravel pits

Time of Flowering: late May through October

Key Distinguishing Characteristic(s): leaves are finely divided (almost carrot-like) and practically odorless when crushed

Impact: chokes out desired vegetation

Plant Shape





Hoary Alyssum – *Berteroa incana*

Also Known As: hoary alison, hoary berteroa, hoary false madwort, hoary false alyssum

General Description: greenish-grey, branched, multi-stemmed perennial with tap root growth; star shaped hairs on seed pods, stems & leaves; in the mustard family

Leaves: ½ to 3 inches (1.3 to 7.6 cm) long, numerous, covered in star-shaped hairs, greenish-grey, smooth edges; lanceolate to oblong shaped; short leaf petioles to no petioles; bottom leaves are basal; alternate leaves on stem

Stems: covered in star-shaped hairs, greenish-grey; one or more stems from the same plant; fine branched tops

Height: up to 36 inches (91 cm) tall

Flower: small white flowers in ball-like clusters; individual flowers 2-3 mm in

diameter with deeply notched petals; numerous

Habitat: sandy or gravelly soils, tame forage, railroad tracks, waste areas, abandoned cropland, roadsides & gravel pits

Time of Flowering: June through September

Key Distinguishing Characteristic(s): star-shaped hairs covering the stem, leaves & seed pods giving the plant a grayish appearance

Impact: competes with other vegetation as it produces seed in high quantities summer-long; taints hay; toxic to horses

Plant Shape





Ox-Eye Daisy – *Chrysanthemum leucanthemum*

Also Known As: white daisy; field daisy; dog daisy, moon penny

General Description: shallow rooted perennial plant with white daisy-like flowers; numerous slightly curving stems from one base; reproduces by seeds (at an average of 859,000 seeds per pound of plant) or by creeping roots

Leaves: lower leaves are dark green, toothed, spoon-shaped and stalked; upper leaves are narrower and stalkless or clasp the stem tightly; leaves are reduced in size moving up the plant; stem leaves are alternate, smooth and glossy

Stems: hairless and erect or slightly curving upwards

Height: 8 to 35 inches (20 to 90 cm) tall; may grow over 40 inches (100 cm) tall in riparian areas

Flower: single daisy-like flowers at ends of stems and main branches; 15 to 30, 0.4 to 0.8 inches (1 - 2 cm)

long white petals on each flower head; centre florets are bright yellow, short, numerous, densely packed, forming a slightly rounded centre; the flower head base has overlapping, light green bracts with brownish margins

Habitat: roadsides, cropland, abandoned cropland, hayland, tame pasture, native prairie & riparian areas

Time of Flowering: early June to late fall

Key Distinguishing Characteristic(s): when crushed, all parts of the plant have a disagreeably sour odour; daisy like flowers; deep green glossy leaves

Impact: very common and often forms dense infestations in pastures, roadsides, riparian areas, gardens and lawns; travels easily along rivers; seeds float.

Plant Shape





Orange Hawkweed – *Hieracium aurantiacum*

Also Known As: devil's weed, devil's paintbrush, orange paintbrush

General Description: perennial with fibrous root system and creeping, leafy stolons; orange, dandelion-like flowers

Leaves: spatula-shaped rosette leaves are 1.5 to 8 inches (3.8 to 20.3 cm) long; bristly hairs cover the leaves; stems are leafless or have small clasping alternate leaves on the lower half

Stems: covered with branched or star-shaped black hairs; one or more stems come from the basal rosette; contain a milky sap

Height: up to 12 inches (30 cm) tall

Flower: yellow-orange to red-orange in colour; 0.5 to 0.75 inches (1.3 to 1.9 cm) wide, dandelion-like with notched petal tips; clustered at the ends of the stems; a single plant can have 1 to 50 flowers

Habitat: tame forage, yard sites, roadsides, gravel pits, native prairie, waste areas and abandoned cropland

Time of Flowering: June through September

Key Distinguishing Characteristic(s): creeping leafy stolons (like strawberries) sprout from the rosette; hairy stems and leaves; yellow-orange to red-orange dandelion-like flowers

Impact: produces a dense mat that chokes out other vegetation

Plant Shape



Allelopathic – the inhibition or retardation of growth in one species of plants by chemicals produced by another species.

Alternate – situated singly (not opposite) on a stem (CPWR).

Auricles – in grasses, the small, ear shaped appendages at the base of a leaf (CPWR).

Awns – a bristle-like appendage at the end of a flower or seed head (CPWR).

Basal – at the base of the plant (CPWR).

Biennial – plants completing their life cycle in two years; usually only flowering in the second year (CPWR).

Bracts – a small reduced/modified scale or leaf often borne below a flower or flower cluster (CPWR/WAP).

Clasping leaves – the leaf base is partially or completely surrounding the stem (CPWR).

Cobwebby – hairs on plant, generally on leaves, resembling the web of a spider in gauziness.

Crown – somewhat enlarged, bottom portion of a plant from which the stem, roots, and/or stolons grow (NRP).

Deeply lobed – a deep rounded projection of a leaf or a leaf-like part of a plant (WAP).

Diffuse – spread out, scattered.

Entire bracts – large, full shaped bracts with smooth edges; without notches or indentations.

Finely divided leaves – leaves that have numerous delicate separations in one leaf; the leaf sometimes appears ‘feather-like’.

Flower head – the grouping of tubular flowers into what appears to be a single flower.

Foliage – the green parts, primarily leaf matter, of a plant.

Glandular hairs – hairs sitting within glands. Glands are small secreting structures that may be protruding, stalked or depressed, and producing oils or nectar (CPWR).

Hybridize – when two plants of a different species within the same genus cross and produce a viable offspring.

Lance-shaped – narrow, but with being wider below the middle and tapering above; generally many times longer than broad (CPWR).

Leaf axil – where the leaf or leaf stem/leaf stalk meets a main stem.

Leaf stalk – joins the leaf to a main stem; also called a leaf stem.

Leaf sheath – the lower part of a leaf blade that clasps and encloses the stem.

Ligule – in grasses, the small, thin appendage projecting from the inner leaf surface at the junction of the stem and the leaf (CPWR).

Glossary

Lobe – a rounded projection of a leaf or a leaf-like part of a plant (WAP).

Mid-rib – the central vein on a leaf (CPWR).

Multi-stemmed – more than one stem arising from the root of a plant.

Nodding – flower head bent downwards.

Nodes – the point of attachment on a stem from which leaves, branches, and in some cases roots arise; often swollen (CPWR).

Oval bracts – bracts in an egg shape with ends equally tapered (WAP).

Perennial – a plant (or part) that lives more than two years (CPWR).

Prostrate – laying flat along the ground.

Rhizomes – a shallow underground stem which stretches out horizontally, producing new shoots and roots along it (CPWR).

Rosette – a circular arrangement of leaves, often on the ground or at the base of the plant (CPWR).

Scalloped – edges with a series of gentle curves.

Spherical – having the shape of a sphere or globe.

Spike – a flower cluster with stalkless individual flowers on a common stalk (WAP).

Spikelets – the combined group of one or more grass flowers attached to the grass stem directly or with a stalk (NRP).

Spur – a hollow projection from the flower petals or centre; usually at the base of a flower (CPWR/WAP).

Stalk – joins the leaf to a main stem; also called a leaf stem.

Stolons – above-ground, horizontal branch from the base of a plant which may take root and produce a new plant (NRP).

Succulent – fleshy or juicy from the accumulation of water in the tissues (CPWR).

Thorns – short, stiff, sharp-pointed appendages; stouter than bristles, prickles, or spines. (CPWR).

Tubular flowers – a tube-shaped grouping of petals, with little expansion outwards. Often, many tubular flowers are grouped together on a flower head (CPWR).

Wing – any thin, flat, usually membranous expansion on a plant part (i.e. wing along a stem) (CPWR).

Glossary definitions obtained from: *Wildflowers Across the Prairies* (WAP); *Common Plants of the Western Rangelands Volume 3 Forbs* (CPWR); *Northern Range Plants* (NRP)

Acknowledgements

Funding for this project has been provided by Agriculture and Agri-Food Canada through the Canadian Agricultural Adaptation Program (CAAP). In Saskatchewan, this program is delivered by the Agriculture Council of Saskatchewan.



Agriculture and
Agri-Food Canada

Agriculture et
Agroalimentaire Canada



Saskatchewan Forage
Seed Development
Commission

Crossing the Medicine
Line Network



This provincial identification guide, *Invasive Plant Species Identification Guide for Saskatchewan*, was developed by the Saskatchewan Forage Council and its partners, and was modeled from the original guide created for Southern Saskatchewan. Funding for this project has been provided by Agriculture and Agri-Food Canada through the Canadian Agricultural Adaptation Program (CAAP). In Saskatchewan, this program is delivered by the Agriculture Council of Saskatchewan.

Funding contributions have also been provided by the Saskatchewan Forage Seed Development Commission, Crossing the Medicine Line Network and True North Specialty Products.

Photos were used with permission from various sources.

Technical review for this guide was provided by project steering committee members including: Julie MacKenzie PAg, Frenchman-Wood River Weed Management Area (WMA); Nancy Johns PAg, Saskatchewan Forage Seed Development Commission; Harvey Anderson PAg, SARM Invasive Plant Species Farm Stewardship Coordinator; John Hauer PAg, Saskatchewan Ministry of Agriculture; Janice Bruynooghe PAg, Saskatchewan Forage Council, and Coy Schellenberg, Saskatchewan Forage Council.

A note of thanks to the Frenchman-Wood River Weed Management Area (WMA), the Native Plant Society of Saskatchewan and previous authors and contributors who laid the groundwork from which this provincial identification guide was adapted.

Acknowledgements

These acknowledgements recognize contributions for the original identification guide:
Invasive Plant Species Identification Guide for Southern Saskatchewan



This identification guide was compiled by Julie MacKenzie P.Ag, Program Coordinator, Frenchman-Wood River Weed Management Area. *The Invasive Plant Species Identification Guide for Southern Saskatchewan* was funded in part by the Saskatchewan Invasive Alien Species in the Community Program and Grasslands National Park and administered by the Frenchman-Wood River Weed Management Area.

Photos were used with permission from various sources. Thank you to Dakota Goddu with Grasslands National Park for collecting these images.

Technical review is an important component of an identification guide. Thank you to Laura James, Grasslands National Park; Sue Michalsky, Crossing the Medicine Line Network; Tara Mulhern Davidson and Kerry Laforge, Agriculture and Agri-Food Canada – PFRA; Garth Wruck and Chet Neufeld, Native Plant Society of Saskatchewan; Clark Brenzil, Saskatchewan Agriculture and Food; Harvey Anderson, SARM/SAF Invasive Alien Plants Group Plan; Gordon Thomas and Julia Leeson, Agriculture and Agri-Food Canada.

Funding through the Invasive Alien Species in the Community Program was provided by the Native Plant Society of Saskatchewan and the Saskatchewan Ministry of Environment through the provincial government's Green Strategy Initiative. It is intended

to assist grassroots collaborative efforts that result in environmental, educational, and economic success against invasive alien species.

The Frenchman-Wood River Weed Management Area (WMA) is a non-profit community based organization that encourages cooperative management across jurisdictional boundaries which are often barriers to invasive species prevention, detection, management, and education. The WMA includes five Rural Municipalities and Grasslands National Park in the south central portion of the province along the US border. The WMA's mission is to work together as landowners, land managers, and local, provincial, and federal governments to manage our invasive and/or noxious weeds, starting with leafy spurge, through prevention, education, early detection, and integrated control measures. The WMA's overarching goal is to protect our local resources and economic stability, riparian areas, native prairie habitat, water quality, natural biodiversity, and aesthetics from invasive and/or noxious weeds.

The Frenchman-Wood River Weed Management Area has been funded in part by the Invasive Alien Species Partnership Program, a Government of Canada initiative, and Grasslands National Park. It is supported by RMs 17, 43, 44, 74 and 75 and the District #3 ADD Board.

Photo Credits:

Front Cover- lft-rt: R. Old, XID Services, bugwood.org; B. Springer, SWA, Swift Current; J. Randall, TNC, tncweeds.ucdavis.edu; T. Korol, Colonsay; USDA, TEAM Leafy Spurge; R. Layson, R. Layson Photography, bugwood.org; C. Evans, River to River CWMA, bugwood.org **Back Cover**- lft-rt: reproduced with permission from *Weeds of the Prairies*, Agdex 640-4, pg. 71, AB Ag & Food; S. Dewey, USU, bugwood.org; UAF Cooperative Extension Archive, U of A, bugwood.org; J. MacKenzie, Hazenmore; C. Evans, River to River CWMA, bugwood.org; R. Holm, U of S, Saskatoon; Texas Weed Information Group, twig.tamu.edu

Pg 1: J. MacKenzie, Hazenmore **Pg 5:** t-b: reproduced with permission from *Weeds of the Prairies*, Agdex 640-4, pg. 2, AB Ag & Food; *Prairie Grasses*, pg. 8, Agriculture Canada, 1971. Reproduced with permission of the Minister of Public Works and Government Services Canada, 2007. **Pg 6:** *Budd's Flora of the Canadian Prairie Provinces*, pg 15, fig. 3 & 4, Agriculture Canada, 1979. Reproduced with permission of the Minister of Public Works and Government Services Canada, 2007. **Pg 8:** lft-rt: S. Dewey, USU, bugwood.org; S. Hurst, USDA, bugwood.org; S. Dewey, USU, bugwood.org **Pg 9:** lft-rt: R. Holm, U of S, Saskatoon; R. Holm, U of S, Saskatoon

Pg 10: lft-rt: R. Old, XID Services, bugwood.org; C. Evans, River to River CWMA, bugwood.org; R. Old, XID Services, bugwood.org **Pg 11:** lft-rt: R. Innes, Powell River BC; S. Dewey, USU, bugwood.org **Pg 12:** lft-rt: USU Archives, USU, bugwood.org; S. Dewey, USU, bugwood.org **Pg 13:** lft-rt: R. Old, XID Services, bugwood.org; W. VanDyk Evans, bugwood.org **Pg 14:** lft-rt: MT Statewide Noxious Weed Awareness & Education Program Archive, MSU, bugwood.org; MT Statewide Noxious Weed Awareness & Education Program Archive, MSU, bugwood.org; N.E. Rees, USDA, bugwood.org **Pg 15:** lft-rt: S. Dewey, USU, bugwood.org; S. Dewey, USU, bugwood.org; S. Dewey, USU, bugwood.org **Pg 16:** lft-rt: S. Dewey, USU, bugwood.org; S. Dewey, USU, bugwood.org; N.L. Jennings, Calgary, eflora.bc.ca **Pg 17:** lft-rt: M. Harte, bugwood.org; S. Dewey, USU, bugwood.org; UAF Cooperative Extension Archive, U of A, bugwood.org **Pg 18:** lft-rt: S. Dewey, USU, bugwood.org; R. Layson, R. Layson Photography, bugwood.org; S. Dewey, USU, bugwood.org **Pg 19:** lft-rt: C. Brenzil, SK Ministry of

Ag, Regina; AIPC, invasiveplants.ab.ca; C. Brenzil, SK Ministry of Ag, Regina

Pg 20: lft-rt: J.D. Byrd, MS-SU, bugwood.org; S. Dewey, USU, bugwood.org; E. Coombs, Oregon Dept. Ag., bugwood.org **Pg 21:** lft-rt: J. MacKenzie, Hazenmore; Richard Old, XID Services, Inc., bugwood.org; Richard Old, XID Services, Inc., bugwood.org **Pg 22:** lft-rt: M.E. Harte, bugwood.org; S. Sweeney, themondaygarden.com; S. Sweeney, themondaygarden.com **Pg 23:** lft-rt: S. Dewey, USU, bugwood.org; S. Dewey, USU, bugwood.org; F. & K. Starr, USGS, bugwood.org **Pg 24:** lft-rt: C. Roche, bugwood.org; reproduced with permission from *Weeds of the Prairies*, Agdex 640-4, pg. 39, AB Ag & Food; C. Roche, bugwood.org **Pg 25:** lft-rt: reproduced with permission from *Weeds of the Prairies*, Agdex 640-4, pg. 37, AB Ag & Food; S. Dewey, bugwood.org; reproduced with permission from *Weeds of the Prairies*, Agdex 640-4, pg. 37, AB Ag & Food **Pg 26:** lft-rt: V. Skilton, eflora.bc.ca; R. Old, XID Services, bugwood.org; V. Skilton, eflora.bc.ca **Pg 27:** lft-rt: G.A. Cooper, USDA, bugwood.org; Texas Weed Information Group, twig.tamu.edu **Pg 28:** lft-rt: reproduced with permission from *Weeds of the Prairies*, Agdex 640-4, pg. 71, AB Ag & Food; J. Riley, eflora.bc.ca; A. Rousseau, eflora.bc.ca **Pg 29:** lft-rt: R. Old, XID Services, bugwood.org; S. Dewey, USU, bugwood.org

Pg 30: lft-rt: C. Evans, River to River CWMA, bugwood.org; MT Statewide Noxious Weed Awareness & Education Program, MSU, bugwood.org; C. Evans, River to River CWMA, bugwood.org **Pg 31:** reproduced with permission from *Weeds of the Prairies*, Agdex 640-4, pg. 101, AB Ag & Food **Pg 32:** lft-rt: N. Grey, SK Invasive Alien Plants Group Plan, Watrous; R. Holm, U of S, Saskatoon; SK Ministry of Ag., Regina **Pg 33:** lft-rt: Catherine Herms, The Ohio State University, bugwood.org; Catherine Herms, The Ohio State University, bugwood.org; Richard Old, XID Services, Inc., bugwood.org **Pg 34:** lft-rt: Mary Ellen (Mel) Harte, bugwood.org; Richard Old, XID Services, Inc., bugwood.org; Steve Dewey, Utah State University, bugwood.org **Pg 35:** lft-rt: M. Shephard, USDA Forest Service, bugwood.org; M. Shephard, USDA Forest Service, bugwood.org; UAF Cooperative Extension Archive, U of A, bugwood.org

References:

- Alberta Agriculture and Food. Restricted, Noxious, and Nuisance Weeds in Alberta. 2007. [www1.agric.gov.ab.ca/\\$department/deptdocs.nsf/all/faq8261](http://www1.agric.gov.ab.ca/$department/deptdocs.nsf/all/faq8261).
- Alberta Invasive Plants Council. Common Tansy, Scentsless Chamomile, Dames Rocket, Field Scabious, & Ox-Eye Daisy fact sheets. 2007. www.invasiveplants.ab.ca.
- Bubar C.J., McColl S.J., & Hall L.M. *Weeds of the Prairies*. Alberta Agriculture and Food. 2000.
- Calliham R.H., Wilson L.M., McCaffrey J.P., & Miller T. *Hawkweeds*. Pacific Northwest Extension. 1997.
- CFIA. Tamarix spp.-Saltcedar, A Canadian Concern. CFIA. 2007.
- Darbyshire S.J. Inventory of Canadian Agricultural weeds. 2003. <http://psd.pwgscc.gc.ca/collection/A42-100-2003E.pdf>.
- E-Flora of BC. Atlas Pages. 2007. www.eflora.bc.ca.
- Government of Canada. Weed Seed Order & Class Definitions. CFIA. 2005. www.justice.gc.ca/en/showfull/doc/cr/SOR-2005-220//en.
- Government of Montana. Montana Noxious Weeds List. 2003. <http://agr.mt.gov/weedpest/noxiousweedslist.asp>.
- Government of Saskatchewan. *The Noxious Weeds Designation Regulations*. Government of Saskatchewan. 1999.
- Hargrave A.N. *Seeded Plants for Forage and Reclamation in Saskatchewan*. Saskatchewan Forage Council. 2007.
- Harris P. & Jensen K.I.N. Round-Leaved Mallow *Malva pusilla* Sm. 2007. http://res2.agr.ca/lethbridge/weedbio/plant/brndmalo_e.htm.
- Invasive.org. Biology and Control of Purple Loosestrife. Dames Rocket fact sheet. 2005. www.invasive.org.
- Jacobs J. & Sing S. Ecology and Management of Dalmation Toadflax (MT3) & Ecology and Management of Yellow Toadflax (MT6). USDA-NRCS. 2006.
- Jacobs J. & Wiese J. Ecology and Management of Invasive Hawkweeds (MT16). USDA-NRCS. 2007.
- Johnson W.S., Wilson R. & Graham J. *Invasive Weed Identification for Nevada*. University of Nevada. 1999.
- King County Noxious Weed Control Program. Hawkweed BMPs. 2005. <http://dnr.metrokc.gov/weeds>.
- Looman J. *Budd's Flora of the Canadian Prairie Provinces*. Agriculture and Agr-Food Canada. 1979.
- Lym R.G. Dalmation Toadflax and Yellow Toadflax Identification and Control. 2002. www.ag.ndsu.edu/pubs/plantsci/weeds/w1239w.htm.
- Manitoba Agriculture, Food and Rural Initiatives. Declaration of Noxious Weeds in Manitoba. 2006. www.gov.mb.ca/agriculture/crops/weeds/fab64s00.html.
- Moss E.H. *Flora of Alberta*. (2nd Ed) Revised by Packer J.G. University of Toronto Press. 2000.
- Parker D. & Reiser M.H. *Low-Impact, Selective Herbicide Application for Control of African Rue*. USDA Forest Service. 1997.
- Saskatchewan Agriculture and Food. Round-Leaved Mallow. 1984. www.agr.gov.sk.ca/docs/production/rndmallw.asp.
- Saskatchewan Agriculture and Food. Scentsless Chamomile, Biology and Management. 2004. www.agr.gov.sk.ca/docs/production/schamomile.asp.
- Stone C. & Lawrence D. *Northern Range Plants*. Alberta Agriculture, Food and Rural Development. 2000.
- Tannas, K. *Common Plants of the Western Rangelands, Volume 3 Forbs*. Olds College & Alberta Agriculture, Food and Rural Development. 2004.
- TNC. Elements of Stewardship Abstract of *Lythrum salicaria*. 2007. www.tncweeds.ucdavis.edu.
- USDA Fire Effects Database. *Bromus japonicus*. 2007. www.fs.fed.us/database/feis/plants/graminoid/brojap/all/html
- USDA Forest Service. Weed of the Week- Cheatgrass & Whitetop, Dame's Rocket & Hoary Alyssum. 2005 & 2006. www.na.fs.fed.us/fhp/invasive_plants.
- Vance F.R., Jowsey J.R., McLean J.S. & Switzer F.A. *Wildflowers Across the Prairies*. Western Producer Prairie Books. 1999.
- Weeds BC. Yellow Starthistle. 2007. www.weedsbc.ca/weed_desc/yel_star.html.
- Wilson L.M., Jette C., Connett J. & McCaffrey, J.P. Biology and Biological Control of Yellow Starthistle. 1998. www.invasive.org/weeds/starthistle/chapter1.html.
- Wright W.H. & Steins I. *Weeds of Canada*. Canada Department of Agriculture. 1970.



For additional copies or information contact:

Saskatchewan Forage Council

306-966-2148

office@saskforage.ca



Funding for this project has been provided by Agriculture and Agri-Food Canada through the Canadian Agricultural Adaptation Program (CAAP). In Saskatchewan, this program is delivered by the Agriculture Council of Saskatchewan.



AGRICULTURE COUNCIL
OF SASKATCHEWAN INC.



Agriculture and
Agri-Food Canada

Agriculture et
Agroalimentaire Canada